

Abstract Submitted
for the DPP95 Meeting of
The American Physical Society

Sorting Category: 4.5 (theoretical)

Nonlinear Laser Filamentation Simulation in 3D* C. H. STILL, R. L. BERGER, A. B. LANGDON, L. V. POWERS, E. A. WILLIAMS, P. E. YOUNG, *Lawrence Livermore National Laboratory* — Recent application of our laser filamentation code to high temperature hohlraums (*e.g.*, $I = 10^{16} \text{W/cm}^2$, $T_e = 10 \text{keV}$), or channeling experiments where nearly all of the mass is evacuated from a cavity, have motivated the development, and integration into **F3d**, of a 3D nonlinear eulerian hydrodynamics (**Nh3**). We have also added a linearized non-local thermal heat conduction model, allowing simulation of thermally driven, as well as ponderomotively driven, filamentation, and a 2nd order wave equation solver. The specifics of **Nh3** and some applications to beam deflection were reported last year.¹ In this presentation, we will show **F3d** simulations for high temperature hohlraums where the filamentation gain per speckle is large, when an extremely tight focus in a plasma is achieved (similar to Peter Young's experiments on Janus)² and in channeling experiments where near vacuum is achieved.

*Work performed under the auspices of the U. S. DoE by LLNL under contract No. W-7405-ENG-48.

¹C. H. Still et al., "3D Nonlinear Hydrodynamics with Beam Deflection Applications", APS/DPP, Louisville KY, 6-11 November, 1995.

²P. E. Young et al., "Laser beam propagation and channel formation in underdense plasmas", *Phys. of Plasmas*, **2** 7 (1995).

☐
☒

Prefer Oral Session
Prefer Poster Session

Charles H. Still
bert@icf.llnl.gov
Lawrence Livermore National Laboratory

Date submitted: July 3, 1996

Electronic form version 1.1